



Research Paper

Ichthyofaunal fresh water biodiversity of Trai belt region district Lakhimpur Kheri Uttar Pradesh, India.

K. C. Yadav*, Dhruv Kumar, Arti Devi, Rajesh Kumar and J. P. Yadav

College of Fisheries Science and Research Centre (Chandra Shekhar Azad University of Agriculture and Technology), Campus, Etawah, Uttar Pradesh, India

*Corresponding author email: kailashyadav447210@gmail.com

Received: 30/06/2026

Revised: 07/07/2026

Accepted: 18/07/2026

Abstract: India is a mega biodiversity hot spot contributing 11.72% of globe fish biodiversity with (2800) species (marine and fresh water). Total fish production has contributed 1.07 % in Gross Domestic Product (GDP) and 5.34% contribution in agricultural GDP. Uttar Pradesh itself have 124 fish spiciness 74 genera and 28 families and contributed about 14.11% Indian fresh water fish diversity. The present study based on the diversity of Lakhimpur Kheri district. Several species are founded during the survey of local area which included in the 10 order, 21 family and 70 species which are distributed among several water bodies located in the district. The study revealed that biodiversity of fishes was declining due to several anthropogenic and natural activities so it's needed to be conserved with several aspects.

Keywords: Fish Biodiversity, Lakhimpur, Fisheries Diversity Conservation.

Introduction:

Fishes are the major part of vertebrate which play important role in animal

biodiversity with 32000 species (marine and fresh water) in global level. India is a mega biodiversity hot spot contributing 11.72% of globe fish biodiversity with (2800) species. Total fish production has contributed 1.07% in Gross Domestic Product (GDP) and 5.34% contribution in agricultural GDP. Uttar Pradesh have 124 fish spiciness 74 genera and 28 families and contributed about 14.11% Indian fresh water fish diversity (Talwar and Jhingran, 1991; Dwivedi et al, 2004, 2007; Kumar et al, 2013; Sarkar et al. 2015). The fish biodiversity of the state is decline rapidly during last few decades due to anthropologic activities, urbanization, environmental changes and degradation, irrigation and power generation and pollution which are directly affected to the natural water bodies and cause direct effects on fish biodiversity (Pandey and Das, 2006; Lakra and Pandey 2009; Lakra 2010; Ayyappan et al, 2011) Lakhimpur Kheri is the largest (in area wise) district of Uttar Pradesh state in India has large aquatic resources for fish production and considerable scope of fresh water fisheries and their development for aquaculture

production. It is an area of Traibelt region and its fisheries resources are distributed in river (Sharda, Ghaghra, Gomti, Gerua, Suheli, Kandwa, Ull, etc.), wetland, lakes; pond and tanks which have billion cubic meter of water every year into main stream. The Ghaghra contributes the largest amount of water per annum to the entire Ganga basin, (94.4 billion cubic meters) (Sarkar et. al. 2015). These water resources contain rich fisheries diversity fishes contain families like Cyprinidae, Bagridae, Siluridae, Mastacembelidae and Schibeidae. Some exotic fish were also affected to the biodiversity of the local fish fauna with their appearance and economic importance among the fish farmers (Singh and Mishra, 2001; Singh et al, 2010; Mayank et al, 2011; Pathak et al, 2011). This district

also has protected forest and wild life areas like Dudhwa Tiger Reserve, Kisanpur sanctuary and Kartaniya Ghkarat wild life sanctuary. That's protected areas also have rich aquatic biodiversity like Fishes, Amphibians' Reptiles and other animals. This fauna was reported by different authors but its required sustainable management. The present study provides baseline database, freshwater fish diversity and their conservation of this area.

Study area:

Different water bodies of district Lakhimpure Kheri is a largest area wise district of Uttar Pradesh state in India. It's situated at the border of Nepal and neighbour district are Bahrich, Sitapur, Hardoi, Shahajhanpur and Piliphit.

Details of Study Sites

Sr. No.	Name of sampling site		Latitude and longitudes of sampling site
1	Sharda Nagar	S-01	28.065523°N 80.965967°E • 151 m
2	Bhira	S-02	28.328927°N 80.495577°E • 184 m
3	Palia	S-03	28.434607°N 80.573629°E • 181 m
4	Majhgayi	S-04	28.346748°N 80.693495°E • 175 m
5	Shrinagar	S-05	28.091843°N 80.791439°E • 162 m
6	Khambhi	S-06	28.098593°N 80.882212°E • 161 m
7	Dhaurahara	S-07	27.999390°N 81.094776°E • 152 m
8	Pachapedi Ghat	S-08	28.127788°N 80.841963°E • 158 m
9	Mahewa Ganj	S-09	27.977224°N 80.785021°E • 159 m
10	Lilauti Nathmandir	S-10	28.008265°N 80.797524°E • 158 m

Material and methods-

Data collection

During study period site selected Sharda Nagar, Bhira, Palia, Majhgayi, Shrinagar, Khambhi, Dhaurahara, Pachapedi Ghat, Mahewa Ganj and Lilauti Nathmandir.

The fishes were collected by the use of various gears and craft like drag net, gill net, purse net, sin net, dip net and fishing boat. The data were collected with the help of local fisherman from different fishing sites and locations of the Tarai belt and

their rivers. This research data was collected during study period of March 2016 to February 2018. For research and data collection several sites and rivers were selected like Ghaghra, Sharda, Suheli, kandwa and Ull.

Fish identification –

The identification of fish specimens were done by the morphological, meristic as well as anatomical character basis. Analysis of morphological traits like body shape, size, colour, lateral line position, lateral line scale, mouth shape size and positions, number of fins, fin size, Fin shape, fin spine, fin rays, position of fins,

body scale (present or absent), barbils (present or absent), number of barbils, rosets, and oculus. Fishes were identified by the help of tools and guidelines from (Day 1878, Talwar and Jhingran (1991), Fishes of Uttar Pradesh and bihar by Gopal Ji Srivastwa, 2022 edition.)

Result and discussion

Several species are founded during the survey of local area during study period which were included in the 09 order, 21 family and 70 species which are distributed in the several water bodies located in the district of the Lakhimpur Kheri, Uttar Pradesh, India.

Fish Distribution and Exotic Species Contribution


Name of Order	Siluriformes	Perciformes	Clupeiformes	Synbranchiformes	Cypriniformes	Mugiliformes	Belontiiformes	Tetraodontiformes	Osteoglossiformes
Number of Species	19	09	01	03	29	02	02	02	02
Number of Exotic Species	02	-	-	-	09	-	-	-	-
Exotic Species Contribution in %	10.53%	-	-	-	31.03%				

Species wise Distribution of Fishes


S. N.	Order	Family	Scientific Name	Common Name / Local Name
1	Siluriformes	Bagridae	<i>Speretaaor</i> (Hamilton, 1822)	Tengan
			<i>Sperata seenghala</i> (Sykes, 1839)	Tengan
			<i>Sperata aur</i>	Aur/ Tengra
			<i>Mystus cavasius</i>	Tengra
			<i>Mystus vitatus</i>	Tengra
			<i>Mystus tengra</i>	Tengra
			<i>Rita rita</i>	Belgegra / ptharchatti
		Siluridae	<i>Ompak pabda</i>	Pabda
			<i>Ompak bimaculatus</i>	Butter catfish
			<i>Wallago attu</i>	Padhin
		Schilbeidae	<i>Alia coilia</i>	Patasi
			<i>Eutropichthy svacha</i>	Vachi
		Sisoridae	<i>Bagarius bagarius</i>	Gonchh
		Clariidae	<i>Clarias batrachus</i>	DesiMangur
			<i>Clarias gariepinus</i> (Exotic cat fish)	Thai mangur
		Heteropneustidae	<i>Heteropneustes fossilis</i>	Singhi
		Pangasiidae	<i>Pangasius pangasius</i>	Pangas
			<i>Pangasiu ssuchi</i>	Pangas
			<i>Pangasius pagasiodon</i> (Exotic cat fish)	Pangas
2	Perciformes	Channidae	<i>Channa punctatus</i>	Girai
			<i>Channa striatus</i>	Sauri
			<i>Channa gachua</i>	Saur
			<i>Channa marulius</i>	Sol
		Ambassidae	<i>Chanda nama</i>	Chanari
			<i>Chanda ranga</i>	Chanari
		Anabantidae	<i>Anabu stetiduneus</i>	Kawai
		Cichlidae	<i>Oreochromis mossambicus</i>	Tilapia
		Nandidae	<i>Nandus nundus</i>	Dhebari
3	Clupeiformes	Clupeidae	<i>Gudusia chapra</i>	Suhiya
4	Synbranchiformes	Mastacembelidae	<i>Mastacembalus armatus</i>	Bam
			<i>Mastacembalus panculus</i>	Bamla
		Synbranchidae	<i>Amphipnouscuchia</i>	Sapayi / Cuchiya eel
5	Cypriniformes	Cyprinidae	<i>Catlacatla</i>	Catla / Bhakur
			<i>Labeo rohita</i>	Rohu
			<i>Labeo calbasu</i>	Karauchh
			<i>Labeo bata</i>	Bata
			<i>Labeo gonius</i>	Kurai

			<i>Bengana dero</i>	Kalabans	
			<i>Labeo angra</i>	Rain	
			<i>Cirrhinus mrigala</i>	Mrigal / Nain	
			<i>Cirrhinus reba</i>	Raiya	
			<i>Hypophthalmichthyes moltrix</i>	Silver carp	
			<i>Cyprinus carpio</i>	Common carp	
			<i>Cyprinus carpio</i> var. <i>specularis</i>	Mirror carp	
			<i>Cyprinus carpio</i> var. <i>nudus</i>	Leather carp	
			<i>Ctenopharyngodon idella</i>	Grass carp	
			<i>Osteobramacotio</i>	Gurda	
			<i>Puntius sarana</i>	Barb	
			<i>Puntius ticto</i>		
			<i>Puntius titius</i>		
			<i>Punteus chola</i>		
			<i>Puntius sphore</i>		
			<i>Amblypharyngodon mola</i>	Dhawai	
			<i>Aristichthys nobilis</i> (Exotic carp)	Bighead carp	
			<i>Hypophthalmichthys molitrix</i> (Exotic carp)	Silver carp	
			<i>Ctenopharyngodon idella</i> (Exotic carp)	Grass carp	
			<i>Cyprinus carpio</i> (Exotic carp)	Common carp	
			<i>Hypophthalmichthys nobilis</i> (Exotic carp)	Bighead carp	
			<i>Colossomama cropomum</i> (Exotic carp)	Pacu carp	
			<i>Oreochromis niloticus</i> (Exotic carp)	Tilapia	
			<i>Oreochromis mossambicus</i> (Exotic carp)	Tilapia	
			Cobitidae	<i>Botia lohachata</i>	Bagha
				<i>Lepidocephalichthys guntea</i>	Nakati
6	Mugiliformes	Mugilidae	<i>Rhinomugil corsula</i>	Corsula	
			<i>Sicamugilca scasia</i>	Mullet	
7	Beloniformes	Belonidae	<i>Xanentodoncancila</i>	Kaua	
			<i>Xenentodon xenentodon</i>	Badakaua	
8	Tetraodonitiformes	Tetraodontidae	<i>Tetraodon cutcutia</i>	Puffer fish/ Fulani	
9	Osteoglossiformes	Notopteridae	<i>Chitala chitala</i>	Moh / moi	
			<i>Notopterus notopterus</i>	Featherback/ Patra	

Graphical Distribution of Species Wise and their % Contribution




Biodiversity Distribution of Family with number of Species



Exotic fish species

S. N.	Order	Family	Scientific name	Common name / local name
	Cypriniformes	Cyprinidae	<i>Cyprinus carpio</i>	Common carp
			<i>Cyprinus carpio</i> var. <i>specularis</i>	Mirror carp
			<i>Cyprinus carpio</i> var. <i>nudus</i>	Leather carp
			<i>Ctenopharyngodon idella</i> (Exotic carp)	Grass carp
			<i>Hypophthalmichthys molitrix</i> (Exotic carp)	Silver carp
			<i>Aristichthys nobilis</i> (Exotic carp)	Bighead carp
			<i>Colossoma macropomum</i> (Exotic carp)	Pacu carp
			<i>Oreochromis niloticus</i> (Exotic carp)	Tilapia
			<i>Oreochromis mossambicus</i> (Exotic carp)	Tilapia
	Siluriformes	Clariidae	<i>Clarias fariatus</i> (Exotic cat fish)	Thai mangur
		Pangasiidae	<i>Pangasius pangasius</i>	Pangas



Exotic contribution in total biodiversity

In exploratory survey total 70 fish species were reported in the fish biodiversity of Lakhimpur Kheri district in similar studies the 61 species were reported by Gupta and Tripathi, 2017 and 56 fish species reported by Sarkar et al. 2010 in Gomti River. These several Species were identified as fresh water species which came from 10 orders and 21 families which included a large part of the Uttar Pradesh fish biodiversity as well as India. During this study the 11 exotic fish species have founded which are occupied in several water bodies but more population near the domestic cultural ponds where there local fish farmer culture fish species for high commercial system.). Sarkar et. al., (2010) similar fish diversity reported in their work on in Gomti river total 70 species, 9 order, 21 families and 42 genera. Singh (2014) suggested and reported in their study that the alien species have potential to disrupt the biological diversity of natural Ecosystem and In the Gomti river 56 fish species belonging 20 families and 42 genera were reported also reported by Srivastava and Singhal (2015). In other similar studies reported that the 56 species belonging to 41 genera, 9 orders and 21 families by Farah Bano & Mohammad Serajuddin (2016).

Conclusion

Based on study and analysis about biodiversity of Lakhimpur Kheri the conclusion drawn the most of species found in least count which indicated the degradable status of the biodiversity similar study reported by Patra and Saha (2013) and Sharma et al. (2014). So it is need to proper action and regulation with challenges for fish faunal diversity conservation through several aspects like the use of various fishing gears and the netting in natural water bodies with proper manners without extra efforts with cause degradation of the biodiversity resources, reduced anthropological activities,

management of natural water resources, shares the benefits of biological resources in an equitable manner and also conserves fish biodiversity through aquaculture production technology.

References

- Talwar, P. K. and Jhingran, A. G. (1991) Inland Fishes of India and Adjacent countries. Vol. 1 & 2, Oxford & IBH Co. Pvt. Ltd. New Delhi.
- Dwivedi, A. C., Tewari, N. P. and Singh, K. R. (2004) Present structure of capture and culture fishery of the Faizabad district (U.P.). Bioved 15, 95-98.
- Dwivedi, A. C., Tewari, N. P. and Mayank, P. (2007) Biodiversity of fishes of Faizabad district (U.P.). Flora & Fauna 13, 379-383.
- Jitendra, K., Pandey, A. K., Dwevedi, A. C. Kumar, A. S. Naik, V., Mahesh, A. S. and Benakappa, (2013) Ichthyofaunal diversity of district Faizabad (Uttar Pradesh). India. J. Exp. Zool. India, 16, 1, 149-154.
- Sarkar, U. K., Dubey, V. K., Pathak, A. K. and Mahapatra, B. K. (2015) Biodiversity of freshwater fishes in Uttar Pradesh: Regenerating status and reinforcing conservation effort for sustainable management. International day for biological diversity biodiversity for sustainable development. Uttar Pradesh state biodiversity board, pp. 88-92.
- Pandey, A. K. and Das, P. (2006) Current status of fish germplasm resources of India and strategies for conservation of endangered species. In: Proceedings of Recent Advances in Applied Zoology (eds. Singh H S, Chaubey A K and Bhardwaj S K), 1-39. Ch. Charan Singh University, Meerut.
- Lakra, W. S. and Pandey, A. K. (2009) Fish germplasm resources of India with special emphasis on conservation and rehabilitation of threatened species. In: Aquaculture Management (eds: Goswami

U C and Kumar D), 85-104. Narendra Pub. House, Delhi.

Lakra, W. S. (2010) Fish biodiversity of Uttar Pradesh: issues of livelihood security, threats and conservation. In: National Conference on Biodiversity, Development and Poverty Alleviation (22th May, 2010), 40-45. Uttar Pradesh State Biodiversity Board, Lucknow.

Ayyappan, S., Moza, U., Gopalakrishnan, A., Meenakumari, B., Jena, J. K. and Pandey, A. K. (2011) Handbook of Fisheries and Aquaculture. 2nd edn. Indian Council of Agricultural Research, New Delhi.

Singh, A. K. and Mishra, A. (2001) Environmental issues of exotic fish culture in Uttar Pradesh. J. Environ. Biol. 22, 205-208.

Singh, A. K., Pathak, A. K. and Lakra, W. S. (2010) Invasion of an exotic fish, common carp, *Cyprinus carpio* L. (Actinopterygi: Cypriniformes: Cyprinidae) in the Ganga River, India and its impacts. Acta Ichthyol. Piscat. 40, 11-19.

Mayank, P., Arvind Kumar and Dwivedi, A. C. (2011) Alien fish species *Oreochromis niloticus* (Linnaeus, 1757) as a powerful invader in the lower stretch of the Yamuna River. Bioved 22, 65-71.

Pathak, R. K., Gopesh, A. and Dwivedi, A. C. (2011) Alien fish species, *Cyprinus carpio* var. *communis* (common carp) as a powerful invader in the Yamuna River at Allahabad, India. Natl. Acad. Sci. Letter 34, 367-373.

Day, F. (1878) Fishes of India, William, Dawson's London U.K. reprint edition, Today and Tomorrow Book agency, Delhi, 1; 2.

Gopal Ji Srivastwa (2022) Fishes of Uttar Pradesh and bihar. 2022 edition.

Gupta, D. and Tripathi, M. (2017) Present status and diversity of ichthyofauna at five selected sites of the Gomti River, Lucknow (India), International Journal of Fauna and biological studies, 4(1): 49-56.

Sarkar, U. K., Gupta, B. K., Lakra, W. S. (2010) Biodiversity, Eco-hydrology, threat status and conservation priority of the freshwater fishes of river Gomti, a tributary of river Ganga (India). Environmentalist. 30:3-17.

Singh, A. K. (2014) Emerging alien species in Indian Aquaculture: prospects and threats, Journal of Aquatic Biology & Fisheries, Vol. 2(1): 32-41.

Srivastava, A., Singhal, A. (2015) Biodiversity, Ecological status and Conservation priority of the fishes of river Gomti, Lucknow (U.P., India). International Journal of Advanced Research. 3(9):1471-1480.

Farah Bano and Mohammad Serajuddin (2016) Biodiversity, Threat Status and Conservation Priority of Ichthyofauna of River Gomti at Lucknow Region, India, Biodiversity Journal, 7 (4): 913-922.

Patra, B. C. and Saha, M. K., (2013) Present Status of Ichthyofaunal Diversity of Damodar River at Burdwan District, West Bengal, India. International Journal of Scientific & Research Publications, 3: 1-11.

Sharma, R., Borana K., Gupta, S. K., Pandey, V. and Khan, A. (2014) Biodiversity and Composition of Ornamental Fish Fauna inhabiting in Upper Lake of Bhopal (M.P.). International Journal of Pharmaceutical and Life Sciences, 5: 3651-3659.